



Visualization of Coronary Wall Atherosclerosis in Asymptomatic Subjects and Patients with Coronary Artery Disease Using Magnetic Resonance Imaging

Citation

Gerretsen, Suzanne C., M. Eline Kooi, Alfons G. Kessels, Simon Schalla, Marcus Katoh, Rob J. van der Geest, Warren J. Manning, et al. 2010. Visualization of Coronary Wall Atherosclerosis in Asymptomatic Subjects and Patients with Coronary Artery Disease Using Magnetic Resonance Imaging. PLoS ONE 5(9): e12998.

Published Version

doi:10.1371/journal.pone.0012998

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:8191190>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Visualization of Coronary Wall Atherosclerosis in Asymptomatic Subjects and Patients with Coronary Artery Disease Using Magnetic Resonance Imaging

Suzanne C. Gerretsen^{1,2}, M. Eline Kooi^{1,2}, Alfons G. Kessels³, Simon Schalla^{2,4}, Marcus Katoh⁵, Rob J. van der Geest⁶, Warren J. Manning⁷, Johannes Waltenberger^{2,4}, Jos M. A. van Engelshoven^{1,2}, Rene M. Botnar⁸, Tim Leiner^{1,2*}

1 Department of Radiology, Maastricht University Medical Center, Maastricht, The Netherlands, **2** Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, The Netherlands, **3** Department of Clinical Epidemiology and Medical Technical Assessment, Maastricht University Medical Center, Maastricht, The Netherlands, **4** Department of Cardiology, Maastricht University Medical Center, Maastricht, The Netherlands, **5** Department of Diagnostic and Interventional Radiology, Saarland University Hospital, Homburg, Germany, **6** Department of Radiology, Division of Image Processing (LKEB), Leiden University Medical Center, Leiden, The Netherlands, **7** Department of Medicine, Cardiovascular Division and Department of Radiology, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts, United States of America, **8** Imaging Sciences Division, King's College, London, United Kingdom

Abstract

Background: Magnetic resonance imaging (MRI) is sensitive to early atherosclerotic changes such as positive remodeling in patients with coronary artery disease (CAD). We assessed prevalence, quality, and extent of coronary atherosclerosis in a group of healthy subjects compared to patients with confirmed CAD.

Methodology: Twenty-two patients with confirmed CAD (15M, 7F, mean age 60.4 ± 10.4 years) and 26 healthy subjects without history of CAD (11M, 15F, mean age 56.1 ± 4.4 years) underwent MRI of the right coronary artery (RCA) and vessel wall (MR-CVW) on a clinical 1.5T MR-scanner. Wall thickness measurements of both groups were compared.

Principal Findings: Stenoses of the RCA (both $<$ and $\geq 50\%$ on CAG) were present in all patients. In 21/22 patients, stenoses detected at MRI corresponded to stenoses detected with conventional angiography. In 19/26 asymptomatic subjects, there was visible luminal narrowing in the MR luminography images. Fourteen of these subjects demonstrated corresponding increase in vessel wall thickness. In 4/26 asymptomatic subjects, vessel wall thickening without luminal narrowing was present. Maximum and mean wall thicknesses in patients were significantly higher (2.16 vs 1.92 mm, and 1.38 vs 1.22 mm, both $p < 0.05$).

Conclusions: In this cohort of middle-aged individuals, both patients with stable angina and angiographically proven coronary artery disease, as well as age-matched asymptomatic subjects, exhibited coronary vessel wall thickening detectable with MR coronary vessel wall imaging. Maximum and mean wall thicknesses were significantly higher in patients. The vast majority of asymptomatic subjects had either positive remodeling without luminal narrowing, or non-significant stenosis.

Trial registration: ClinicalTrials.gov NCT00456950

Citation: Gerretsen SC, Kooi ME, Kessels AG, Schalla S, Katoh M, et al. (2010) Visualization of Coronary Wall Atherosclerosis in Asymptomatic Subjects and Patients with Coronary Artery Disease Using Magnetic Resonance Imaging. PLoS ONE 5(9): e12998. doi:10.1371/journal.pone.0012998

Editor: Alma Zernecke, Universität Würzburg, Germany

Received: April 26, 2010; **Accepted:** August 31, 2010; **Published:** September 29, 2010

This is an open-access article distributed under the terms of the Creative Commons Public Domain declaration which stipulates that, once placed in the public domain, this work may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose.

Funding: This work was supported by the Netherlands Organization for Scientific Research (http://www.nwo.nl/nwohome.nsf/pages/SPPD_5R2QE7_Eng; NWO VENI scheme, Grant number 916.46.034). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: timleiner@gmail.com

† Current address: Department of Radiology, Utrecht University Medical Center, Utrecht, The Netherlands

Introduction

Coronary artery disease (CAD) remains the leading cause of mortality and morbidity in the Western World and developing countries despite continued improvements in prevention and early diagnosis[1,2]. The most frequent cause underlying an acute coronary event is disruption of an atherosclerotic plaque[3,4]. Rupture-prone plaques are referred to as thin-cap fibroatheroma

and characterized pathologically as having a large necrotic core, high macrophage content and a thin, fibrous cap[3,4].

In asymptomatic individuals acute myocardial infarction and sudden death may be the first clinical manifestation of coronary atherosclerosis[5]. About 50–64% of all sudden cardiac deaths occur without prior recognition of coronary heart disease[1]. Well-established risk factors for coronary atherosclerosis are hypercholesterolaemia, hypertension, cigarette smoking, diabetes mellitus,

and systemic inflammation[6,7]. However, despite their usefulness for global risk assessment, these parameters lack specificity for prediction of individual coronary plaque burden.

Detection of subclinical coronary atherosclerosis in high-risk patients might become an important strategy to prevent clinical coronary heart disease. Compared to other imaging modalities magnetic resonance imaging (MRI) has superior ability to differentiate soft tissues and has shown to facilitate characterization of atherosclerotic plaque components in the aorta and carotid artery by using multisequence MRI[8,9,10]. These properties make it uniquely suited for serial non-invasive imaging of the vessel wall, and for monitoring effects of pharmaceuticals on plaque progression or regression. Prior work has demonstrated the capability of MRI to visualize the coronary vessel wall (CVW) with high spatial resolution, as well as vessel wall thickening and positive Glagov-type remodeling in patients[11,12,13].

In this study we sought to evaluate the prevalence and extent of coronary atherosclerosis in a cohort of middle-aged subjects without clinical CAD. MR coronary vessel wall (MR-CVW) characteristics were compared to a positive control group of patients with angiographically confirmed CAD.

Materials and Methods

Subjects

Between October 2005 and February 2007, 26 middle-aged subjects without previous history or clinical symptoms of CAD and 25 patients with symptoms of stable angina and angiographically confirmed CAD were enrolled. Healthy subjects were recruited via newspaper advertisement. Patients were recruited from the cardiology outpatient clinic of our hospital after having x-ray coronary angiography (CAG). Patients with confirmed CAD underwent MRI prior to percutaneous coronary intervention. Hemodynamically unstable patients or patients with previous interventions in the right coronary artery (RCA) as well as patients with severe arrhythmia or contra-indications for MRI were excluded. Approval of the Maastricht University Medical Center review board and written informed consent of all participants were obtained prior to inclusion.

In addition to registration of demographic data and risk factors, the estimated glomerular filtration rate (eGFR) and PROCAM-score[14] were calculated whenever data were available.

MR Imaging protocol

All MR studies were performed on a clinical 1.5T system (Intera, Release 11.1 Philips Medical Systems, Best, The Netherlands) using a dedicated 5-element phased-array cardiac radiofrequency (RF) coil. Subjects were examined in the supine position. First, localizer scans were obtained and the subject specific mid-diastolic trigger delay and acquisition window were determined. These scans were followed by a double oblique oriented three-dimensional (3D) bright blood balanced steady state free precession (bSSFP) coronary MR angiography (CMRA) of the RCA (TR/TE/FA: 6.2/3.1/120°, resolution: 0.98×0.98×3 mm). In the same orientation, a vessel wall scan was acquired (3D FFE, radial k-space sampling, modified double inversion recovery (DIR) prepulse) as previously described[13,15,16]. Imaging parameters for the vessel wall scan were: TR: 8 ms, repeated every other heartbeat for good blood suppression, TE: 2.0 ms, Flip Angle: 30°, Field of view: 300×300 mm and matrix: 384×384. Twenty interpolated slices of 1 mm were obtained. The resulting (acquired) spatial resolution was 0.78×0.78×2 mm. Cardiac gating and a 2D navigator beam were used for compensation of cardiac and respiratory motion. This protocol has previously been

shown to result in highly reproducible images of the coronary vessel wall in healthy volunteers[16]. No contrast agents or sublingual nitroglycerin were used.

Evaluation of coronary artery stenosis and plaque burden

Image quality (IQ) of the vessel wall datasets was scored by two experienced, independent blinded observers on a 4 point scale: 1 = well-defined vessel wall borders and high vessel wall-to-background contrast; 2 = some blurring of the vessel wall borders, average to good vessel-wall to background contrast; 3 = major artifacts or severe blurring of the vessel wall borders, low to average vessel wall-to-background contrast, and 4 = vessel wall cannot be identified, very low vessel wall-to-background contrast (modified from Zhang et al. and Spuentrup et al.[17,18]). An experienced cardiologist blinded to the results of the vessel wall scans assessed the location and degree of a stenosis on the CMRA images. Coronary artery luminograms were classified as negative when there was no visible sign of disease or mild to moderate luminal narrowing (stenosis <50%). Significant disease (stenosis ≥50%) was considered to be present in cases of obvious narrowing of the vessel or marked attenuation of coronary lumen signal[19]. X-ray coronary angiography (CAG) - available in all CAD patients - was visually interpreted by different cardiologists prior to MRI.

Subsequently, an observer blinded to clinical data quantitatively analyzed coronary artery wall images. Minimum, mean and maximum wall thickness were measured on the 3D vessel wall sequence. Lumen diameter was measured on bSSFP images. For measurements, a custom made software package was used (VesselMASS, Leiden University Medical Center, department of Radiology, division Image Processing, Leiden, the Netherlands)[20], adapted for viewing and analysis of longitudinal coronary vessel wall images. For analysis purposes the RCA vessel wall was divided into three segments according to the American Heart Association classification[21].

To prevent errors in wall thickness and lumen diameter measurements due to reconstruction errors, only source images were used for measurements. In source images where the vessel wall was visible, the inner and outer borders of the anterior and posterior vessel wall were manually delineated (Figure 1). Each resulting region of interest (ROI) was automatically divided in 100 subsegments for quantitative analysis of wall thickness. Because measurements in adjacent subsegments are not independent due to partial volume effects we corrected for the length of the measured segment and spatial resolution by only using subsegments from separate voxels and discarding measurements in between. The number of included segments equals the number of voxels in the ROI, which was calculated by dividing the length of the ROI by the in-plane spatial resolution. Coronary plaque burden was quantified as the mean and maximum RCA vessel wall thickness along the proximal two vessel segments[19]. In addition to measuring wall thickness, a normalized wall index was calculated (100*(diameter lumen divided by the sum of anterior and posterior wall thickness)) to get a better quantitative impression about the degree of positive remodeling.

Based on both CMRA and MR-CVW, areas could be classified as diseased or not-diseased according to the following criteria: ROIs with vessel wall thickening were classified as diseased. This could either be in areas of stenosis (combination of luminal narrowing with vessel wall thickening) or in areas with positive remodeling (wall thickening without luminal narrowing). ROIs with normal lumen diameter compared to the rest of the vessel and with a vessel wall that had no areas of focal wall thickening or focal high signal intensity, were classified as "not diseased". Since




Figure 1. Measurement of vessel wall thickness and signal intensity on source images. The anterior and posterior vessel walls are delineated (A). In each segment 100 measurements are made of the thickness and signal intensity of the proximal and distal RCA vessel wall and the diameter of the lumen (B,C). The custom made software program calculates wall thickness, signal intensity, and lumen diameter (D-F) over the entire length of the measured segment.
doi:10.1371/journal.pone.0012998.g001

calcified plaques demonstrate low signal intensity on all MR sequences and can therefore be difficult to visualize, ROIs with obvious luminal narrowing but without visible vessel wall thickening were interpreted as diseased with (heavily) calcified plaque if overall image quality of the vessel wall scan was good and if the vessel wall and lumen could be distinguished in other segments of the vessel.

Statistical analysis

Statistical analysis was performed using SPSS 11.5 (SPSS Inc., Chicago, IL, USA). Differences in IQ between the patient and control group were compared using a Wilcoxon rank-sum test. Minimum, maximum and mean wall thickness and the normalized wall index in both groups were compared using an unpaired student t-test. Differences between men and women in both groups were analyzed using a Wilcoxon rank-sum test.

In addition, we pooled the results by testing the global hypothesis that all results of the measurements (minimum, maximum and mean values) of vessel wall thickness were equal using the ordinary least square test [22,23]. Significance was assumed at two-tailed p values of less than 0.05.

Results

Subject characteristics

All 51 subjects underwent coronary MRI without complications. In three patients, the coronary vessel wall sequence could not be performed due to severe back pain (1 patient) and software problems (2 patients). This resulted in evaluable vessel wall scans in 26 asymptomatic subjects (11 M, mean age 56 ± 4 yrs) and 22 patients (15 M, mean age 60 ± 10 yrs).

Clinical characteristics of this study cohort are listed in table 1. There was a significant difference in subjects' weight and the presence of hypercholesterolemia and hypertension between both groups. In the patient group, there was a trend towards higher prevalence of males and higher age, higher incidence of diabetes, and higher PROCAM score. In patients ($n=12$) in whom a PROCAM score could be calculated, a 10-year risk for acute coronary events of 12.0% (± 8.1 , range: 1.3–25.1%) was present, versus 4.8% (± 2.4 , range: 1.9–8%) in control subjects ($n=6$). Linear regression showed a weak correlation between eGFR and mean wall thickness ($R: 0.54$, $p < 0.01$). In subjects with low eGFR decreased vessel wall thickness was found.

Table 1. Subject characteristics.

	Patients (N = 22)	Asymptomatic subjects (N = 26)	P
characteristics			
Gender, male n (%)	15 (68)	11 (42)	0.08
Age, y (\pm SD)	60.41 (10.35)	56.08 (4.43)	0.08
Weight, kg (\pm SD)	81.27 (15.59)	71.35 (13.73)	0.02
Systolic blood pressure, mm Hg (\pm SD)	137 (14.16)	133 (16.38)	0.39
Diastolic blood pressure, mm Hg (\pm SD)	80 (9.28)	79 (11.46)	0.87
Diabetes, n (%)	3 (13.6)	0 (0)	0.08
Current smoker, n (%)	3 (13.6)	5 (19.2)	0.61
Hypertension, n (%)	13 (59.1)	1 (3.8)	<0.01
Hypercholesterolemia, n (%)	11 (50)	2 (7.7)	0.01
eGFR (\pm SD) mL*min ⁻² * 1.73 m ⁻²	88.61 (27.5) (n = 22)	76.01 (24.2) (n = 9)	0.24
PROCAM-score (\pm SD) [†]	44.6 (9.7) (n = 12)	36.0 (6.0) (n = 6)	0.07

*eGFR: estimated Glomerular Filtration Rate.

[†]PROCAM-score: based on the PROspective Cardiovascular Muenster study-scoring scheme.

doi:10.1371/journal.pone.0012998.t001

Coronary artery atherosclerosis

The prevalence of coronary stenoses as identified by CMRA in the 48 subjects with complete scans is listed in Table 2. Significant coronary artery stenoses ($\geq 50\%$) were found in 17 patients on CMRA, but only in 16 patients on CAG. In 19/26 asymptomatic subjects (73%), there was evidence of at least some luminal narrowing on CMRA. Nine of these 19 asymptomatic subjects had a significant stenosis ($\geq 50\%$). In the patient group, the degree of stenosis as estimated on CMRA ($<$ or \geq than 50%) was compared to the results of CAG. In 21 out of 22 patients (95%), location of stenoses detected on CMRA corresponded to stenoses detected with CAG, although in some segments there was overestimation of the degree of stenosis on CMRA. Agreement between CAG and CMRA for detection of lesions is demonstrated in table 3. Sensitivity and specificity were 100% and 83%, respectively.

Lumen diameter as measured on MR-CVW, was slightly smaller compared to diameters measured on CMRA. In patients, lumen diameter on CMRA was 1.86 ± 0.46 mm, versus 1.66 ± 0.53 mm on MR-CVW ($p = 0.02$). In asymptomatic subjects, lumen diameter was 2.01 ± 0.55 on CMRA and 1.64 ± 0.63 on MR-CVW ($p < 0.01$). The slight difference in in-plane spatial resolution between both techniques (0.99 mm on CMRA,

Table 2. Stenoses on coronary MRA.

	Patients (N = 22)	Asymptomatic Subjects (N = 26)
No stenoses	0 (0)	7 (27)
<50% stenoses	5(23)	10 (38)
$\geq 50\%$ stenoses	17 (77)	9 (35)

Prevalence of stenoses (number (%)) in the right coronary artery as seen with coronary MRA.

doi:10.1371/journal.pone.0012998.t002

Table 3. Comparison between CMRA and CAG on segment level.

		CAG		Non-evaluable segments
		0–50%	$\geq 50\%$	
CMRA proximal	<50%	10	0	0
	$\geq 50\%$	2	10	
CMRA mid (*)	<50%	10	0	1 (CAG)
	$\geq 50\%$	3	8	
CMRA distal	<50%	10	0	4 (CMRA) 1 (CAG)
	$\geq 50\%$	5	2	

Agreement between coronary MRA (CMRA) and X-ray coronary angiography (CAG) on degree of stenoses in the right coronary artery in patients with coronary artery disease (N = 22) in the proximal, mid and distal RCA. Values represent number of segments.

doi:10.1371/journal.pone.0012998.t003

0.78 mm on MR-CVW), can theoretically lead to overestimation of the lumen diameter on CMRA, but also overestimation of coronary stenoses, or underestimation of lumen diameter on MR-CVW. However, on MR-CVW partial volume may play a more important role since the lumen is surrounded by vessel wall. It is more probable that lumen measurements are more accurate on CMRA, although a slight influence of the signal of the vessel wall on diameter measurements on CMRA cannot be ruled out completely. In theory however, we expect that this effect is minimal due to the technique used for suppression of myocardial signal (T2prep). Since the vessel wall is mainly composed of smooth muscle cells, collagen and elastin, the T2 is expected to be similar to the T2 of myocardium. Because of signal suppression due to T2prep, only little signal contribution of the vessel wall can be expected.

There was a trend towards smaller minimum lumen diameter as measured on the luminography sequence in the patient group (1.06 ± 0.58 in patients vs 1.35 ± 0.53 mm in asymptomatic subjects, $p = 0.08$). However, no significant differences could be demonstrated between mean (1.86 ± 0.46 mm in patients, 2.0 ± 0.55 mm in asymptomatic subjects, $p = 0.35$) and maximum (2.91 ± 0.57 mm in patients, 2.73 ± 0.70 mm in subjects, $p = 0.35$) luminal diameters of the two groups (Table 4).

IQ of the vessel wall scans was good in the majority of cases with a mean score of 1.42 ± 0.58 for all subjects. There was no significant difference in IQ between patients and asymptomatic subjects (1.32 ± 0.57 vs 1.5 ± 0.58 for observer 1, $p = 0.21$, 1.42 ± 0.58 vs 1.41 ± 0.59 for observer 2, $p = 0.9$).

Various degrees of non-uniform vessel wall thickening were seen on the MR-CVW scans in patients and asymptomatic subjects (Figures 2 and 3). Quantitative results are summarized in Table 4. Maximum and mean wall thicknesses were significantly larger in the patient group. The normalized wall index was higher in asymptomatic subjects when compared to patients (82.8 ± 20.7 versus 68.2 ± 17.0 , $p = 0.012$), representing an increased wall thickness in all patients and asymptomatic subjects but a relatively thicker vessel wall compared to lumen diameter in patients. Pooled results of wall thickness also demonstrated a significant difference between the groups ($p = 0.03$).

Overall, there was good agreement between luminal narrowing at CMRA and increased wall thickness at MR-CVW in all patients. Agreement between CMRA and MR-CVW in asymptomatic subjects is demonstrated in Table 5. There were 5 asymptomatic subjects with stenoses without wall thickening: in

Table 4. Vessel wall thickness and lumen diameter measurements.

	Patients (N = 22)	Asymptomatic subjects (N = 26)	p
Vessel wall:			
Minimal vessel wall thickness (mm) (±SD)	0.71 (0.19)	0.65 (0.16)	0.269
Maximal vessel wall thickness (mm) (±SD)	2.16 (0.37)	1.92 (0.44)	0.048
Mean vessel wall thickness (mm) (±SD)	1.38 (0.18)	1.22 (0.22)	0.011
Overall (standardized) wall thickness (±SD)	0.87 (2.25)	−0.74 (2.57)	0.026
Lumen CMRA			
Minimal lumen diameter (mm) (±SD)	1.06 (0.58)	1.35 (0.53)	0.080
Maximal lumen diameter (mm) (±SD)	2.91 (0.57)	2.73 (0.70)	0.354
Mean lumen diameter (mm) (±SD)	1.86 (0.46)	2.01 (0.55)	0.346
Lumen MR-CVW			
Minimal lumen diameter (mm) (±SD)	0.91 (0.58)	1.02 (0.53)	0.475
Maximal lumen diameter (mm) (±SD)	2.57 (0.66)	2.39 (0.81)	0.396
Mean lumen diameter (mm) (±SD)	1.66 (0.53)	1.64 (0.63)	0.930
Normalized wall index (%), (±SD)	68.2 (17.0)	82.8 (20.7)	0.012

Vessel wall thickness and lumen diameter measurements (±SD) measured on both CMRA and MR-CVW in patients and healthy volunteers. Normalized wall index was calculated as a value to represent positive remodeling according to the following formula: $100 \times (\text{lumen diameter} / \text{sum of anterior and posterior wall thickness})$.

doi:10.1371/journal.pone.0012998.t004

three of these subjects, the area of stenosis was not well visualized on the vessel wall scan, in one subject vessel wall and lumen could hardly be distinguished and in one subject the location of the vessel wall thickening did not correspond to the area of stenosis on CMRA.

The groups were age-matched, but not matched for gender. There was a trend towards a higher prevalence of women in the group of healthy volunteers (Table 1). In patients, there were no significant differences between men and women. However, in the group of healthy volunteers, there was a significant difference in mean vessel wall thickness, mean luminal diameter on CMRA, mean luminal diameter on MR-CVW and a trend towards difference in normalized wall index (Table 6).

Discussion

This is one of the few studies to use non-invasive MRI to assess coronary artery plaque burden in an asymptomatic population-based cohort of adults using magnetic resonance CVW imaging. The most significant finding of this study is the high prevalence of coronary wall plaque in asymptomatic adults free of known CAD. Four out of 26 healthy subjects exhibited evidence of positive remodeling without luminal narrowing and the majority of subjects (19/26) exhibited evidence of luminal narrowing at CMRA, in some cases even exceeding 50%. As expected, the burden of coronary atherosclerosis was higher in the positive control group, which consisted of patients with angiographically

confirmed CAD. Despite the lack of a gold standard for visualization of the coronary vessel wall (i.e. intravascular ultrasound), these data suggest that MRI may help to non-invasively identify not only late stages of atherosclerosis but also the early manifestations of atherosclerotic disease in the coronary vessel wall.

Our results corroborate the findings of Kim et al, who investigated subjects with longstanding diabetes type 1 but without cardiovascular symptoms[19]. In these asymptomatic patients, only 10% of patients with nephropathy and 0% of patients with normoalbuminuria had evidence of coronary stenoses although the prevalence of visually identifiable coronary plaque in the RCA was higher (76% in patients with nephropathy versus 15% in patients with normoalbuminuria). The prevalence of vessel wall thickening in the RCA is in the same range as in our study group, however, we found more cases of luminal narrowing, probably reflecting differences in study populations. Although diabetic subjects tend to have more atherosclerosis, our asymptomatic subjects were older than the subjects with diabetic nephropathy from the study by Kim et al (mean age of 56 vs 48 years). Additionally, MRI is known to overestimate stenosis and possible slight differences in spatial resolution between the used techniques can contribute to differences in assessment of stenosis. Tuzcu et al[24] studied a population of 262 subjects and found vessel wall thickening of more than 0.5 mm on intravascular ultrasonography (IVUS) in >70% of asymptomatic subjects by the age of 40, while only 8% of X-ray coronary angiograms of the entire cohort demonstrated mild luminal irregularities. While the study by Tuzcu once again underscores that early vessel wall abnormalities go undetected at coronary angiography, there was a large difference in prevalence of luminal irregularities compared to our study. The fact that our study demonstrated luminal irregularities on CMRA in >70% of subjects compared to 8% in the study by Tuzcu et al. can be explained by the better spatial resolution of X-ray coronary angiography compared to CMRA, the fact that flow disturbances on CMRA can lead to overestimation of the degree of stenoses, and the difference in age and case-mix between the populations.

A recent publication from the Multi-Ethnic Study of Atherosclerosis (MESA) also found positive remodeling in a large cohort of 179 subjects without known coronary artery disease[25]. In contrast to our study, the MESA study did not find any significant stenoses ($\geq 50\%$), despite the use of similar spatial resolution as well as an identical technique for suppressing blood flow[25]. Axial images of all three coronary arteries (LM, RCA and LAD) were acquired, but due to time constraints this was limited to the proximal parts of the vessels of interest, with only one slice in the LM and 3 slices in the RCA and LAD. Axial imaging of the coronary vessel wall is very time consuming, and the longitudinal technique that we used offers an overview of the vessel wall in a much larger part of the coronary artery of interest. The difference in acquisition approach, especially the chosen imaging plane and resulting differences in partial volume artefacts, between our study and the MESA study is one of the likely explanations for the discordant results.

Clinical CAD has a long asymptomatic period during which biological risk factors interact with genetic and environmental influences to initiate and promote the development of coronary atherosclerotic plaque[26,27]. Once established, atherosclerosis can exist in a subclinical state characterized by the presence of plaque but an absence of clinical signs and symptoms. Acute clinical manifestations of coronary atherosclerosis such as unstable angina and myocardial infarction are usually the consequence of acute rupture of the atherosclerotic plaque, leading to exposure of




Figure 2. MRA and MR-CVW in asymptomatic subjects. Balanced steady state free precession (bSSFP) coronary MRA (A,C) and corresponding vessel wall scans (B,D) of the right coronary artery (RCA) in asymptomatic subjects. A and B represent the RCA of a 62 y/o female (A,B). There is a thin vessel wall with uniform signal intensity (arrows in B). In C and D, the RCA of a 54 y/o female is shown. In this subject, the posterior proximal vessel wall is thicker and has higher signal intensity compared to the anterior vessel wall (arrows in D), indicating the presence of positive remodeling. Ao indicates aorta.

doi:10.1371/journal.pone.0012998.g002

thrombogenic components to the bloodstream, with superimposed thrombus formation[28]. Although traditional risk factors and risk scores such as the Framingham[7,29] and PROCAM[14] scores are accurate at predicting risk to experience a cardiovascular event in the near future for groups of patients, they often lack discriminative power to characterize individual subjects risk. Approximately 40% of the adult population in the U.S. and Western Europe are at intermediate (0.5–2.0%) and 10% at high risk (>15%) for developing an acute coronary event over the next decade[30]. Therefore, further risk stratification of individual patients with CAD using novel biochemical markers as well as noninvasive imaging techniques may be important. Our study was motivated by the paucity of data on the prevalence and characteristics of early CAD and coronary atherosclerotic plaque burden at MRI in an unselected cohort of asymptomatic middle aged adults free of known CAD at baseline. However, it is known that the measured degree of coronary stenosis that was taken as significant in this study (i.e. $\geq 50\%$) often is not associated with significant impairment in coronary blood flow. This might be the reason why none of the control subjects with stenoses $\geq 50\%$ at MRI were symptomatic. An alternative explanation could be that with MRI luminal irregularities are erroneously overestimated[31,32,33], which is one of the shortcomings of MRI in comparison to conventional X ray angiography.

Whether MR-CVW confers additional information over more established non-invasive imaging techniques such as calcium

scoring[34,35] and multislice computed tomography (MSCT) [36,37] is unknown. Spatial resolution of CT is better. However, in comparison to epicardial coronary artery calcium scoring, MRI can detect earlier changes such as outward remodeling. Outward remodeling has been shown to be associated with plaques that have a higher relative content of fibrofatty components as opposed to calcification[38], as well as with plaques that ultimately exhibit an unstable clinical presentation[39]. The excellent soft-tissue resolution of MRI enables better characterization of plaque components than CT. Furthermore, no contrast agents are needed to visualize the coronary vessel wall with MRI, which is beneficial in patients with renal failure or allergies. An additional advantage of MRI is that it does not require radiation exposure, an issue that takes on considerable importance when weighing risks and benefits in asymptomatic subjects, and when follow-up studies are desired[40].

Limitations of this study should be considered. Due to ethical considerations, no comparison to a standard of reference was obtained in asymptomatic volunteers and therefore, there is no certainty about actual prevalence of abnormalities in this group. Furthermore, the groups were matched for age but not for gender. It is known that there are gender related differences in coronary artery disease. In general, women have a less typical clinical presentation and often have fewer angiographic high-risk features[41,42]. In our group of asymptomatic volunteers, there were several significant differences between men and women.




Figure 3. MRA and MR-CVW in patients with stable angina. X-ray coronary angiography (A,D), balanced steady state free precession (bSSFP) coronary MRA (B,E) and corresponding vessel wall scans (C,F) of the right coronary artery (RCA) in two patients with stable angina. In a 40 y/o male (A–C), vessel wall thickness and signal intensity are increased in areas of stenosis and wall irregularities (arrows). In a 67 y/o female (D–F) the culprit lesion was located in the left anterior descending artery (not imaged). However, the proximal RCA demonstrated wall focal luminal irregularities (arrows in D,E). In F, corresponding focal heterogeneity in wall thickness and signal intensity of the entire vessel wall are clearly visible. Ao indicates aorta.
doi:10.1371/journal.pone.0012998.g003

Women had thinner vessel walls and smaller lumen diameter, however, the lower normalized wall index in women indicates a relatively thick vessel wall compared to lumen diameter, which could be related to more positive remodeling. In patients, these differences between men and women were not found. Our results are supported by the study by Miao et al. who also found gender related differences in lumen size, vessel wall area and mean vessel wall thickness in a large cohort of asymptomatic men and women who underwent MRI of the coronary vessel wall [25]. However, in both men and women, they found a similar, significant,

relationship between vessel wall thickness and outer lumen area or lumen area, both measures for the degree of positive remodeling. In the study by Miao et al, there was no comparison between patients with CAD and healthy subjects. Due to the difference in the amount of women between both groups in our study, future

Table 5. Agreement between MR coronary vessel wall imaging (MR-CVW) and coronary magnetic resonance imaging (CMRA) for detection of stenoses and vessel wall thickening in asymptomatic volunteers.

		CMRA	
		No stenoses	Stenoses
MR-CVW	No Wall thickening	3	5
	Wall thickening	4	14

Values represent number of subjects.
doi:10.1371/journal.pone.0012998.t005

Table 6. Gender related differences in the group of patients (n = 22) and the group of healthy volunteers (n = 26).

	Men	Women	p
<i>Patients (15M, 7F)</i>			
Mean vessel wall thickness (mm) (\pm SD)	1.41 (0.18)	1.32 (0.18)	0.341
Lumen diameter on CMRA (mm) (\pm SD)	1.90 (0.38)	1.79 (0.61)	0.307
Lumen diameter on MR-CVW (mm) (\pm SD)	1.66 (0.49)	1.65 (0.63)	0.805
Normalized wall index (%), (\pm SD)	67.8 (13.0)	69.1 (24.9)	0.916
<i>Healthy subjects (11M, 15F)</i>			
Mean vessel wall thickness (mm) (\pm SD)	1.34 (0.24)	1.14 (0.18)	0.046
Lumen diameter on CMRA (mm) (\pm SD)	2.43 (0.57)	1.69 (0.26)	0.001
Lumen diameter on MR-CVW (mm) (\pm SD)	2.04 (0.75)	1.36 (0.31)	0.005
Normalized wall index (%), (\pm SD)	94.0 (26.6)	74.6 (9.3)	0.092

doi:10.1371/journal.pone.0012998.t006

studies should not only match for age but also for gender to further study possible gender-related differences.

Currently, there are still some limitations in scan technique for coronary vessel wall imaging. With currently used in-plane spatial resolution of 0.78 mm, detection of small variations in wall thickness due to partial volume effects are not yet possible. Additionally, due to the current spatial resolution, it can be expected that partial volume effects account for overestimation of coronary vessel wall thickness. Other factors that lead to overestimation of wall thickness are curvature of the vessel and out of plane orientation of the vessel. However, these factors play a more important role with multiplanar reformatting, and to minimize errors due to these factors, measurements were performed on source images.

Furthermore, because the obvious long-term goal of MR-CVW imaging is not only measurement of plaque size but also composition, increased spatial resolution is needed to characterize plaque components. This is important, as it is known that a large lipid-rich necrotic core and a thin fibrous cap portend a poor prognosis. First attempts at characterization of coronary plaques using contrast-enhanced MR-CVW imaging have been made by Maintz et al[43] and Yeon et al[44], who demonstrated that use of contrast-enhanced MR CVW imaging allowed identification of areas of delayed enhancement that correlated with severity of atherosclerosis by MSCT and quantitative X-ray angiography. Also, whether therapeutic intervention guided by imaging modalities will influence clinical event rates remains an important focus of research. Imaging cannot be regarded as an isolated tool to

determine prognosis. However, it is highly likely that imaging will only confer additional information in selected subgroups of subjects in combination with traditional risk factors and (novel) serological markers. This concept is also known as the 'vulnerable patient'[45]. Further limitations include the relatively low sample size and, because of an attempt to limit total scan time in our subjects, analysis of coronary plaque burden was restricted to the RCA.

In conclusion, although there was low prevalence of significant coronary stenoses, we found coronary atherosclerotic plaque in a large proportion of asymptomatic middle-aged subjects at MR CVW imaging. As expected, there were less stenoses and asymptomatic subjects exhibited lower mean coronary vessel wall thickness compared to a control group of patients with angiographically proven coronary artery disease. Because the time-course from subclinical disease to clinical events is highly variable, it is important that future studies focus on the association between MR CVW evidence of disease and clinically manifest incident CAD.

Author Contributions

Conceived and designed the experiments: WJM JW JMAvE RMB TL. Performed the experiments: SCG. Analyzed the data: SCG MEK AGK SS TL. Contributed reagents/materials/analysis tools: RJvdG. Wrote the paper: SCG MEK AGK SS MK RJvdG WJM JW JMAvE RMB TL. Data interpretation and supervision: MEK JvE TL. Statistical analysis: AK. Assistance with data acquisition in early phase of study: MK. Developed MR method: RB. Handled funding: TL.

References

- Rosamond W, Flegal K, Furie K, Go A, Greenlund K, et al. (2008) Heart disease and stroke statistics—2008 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 117: e25–146.
- Mackay J, Mensah GA (2004) The atlas of heart disease and stroke. Geneva: World Health Organization. 112 p.
- Burke AP, Farb A, Malcom GT, Liang YH, Smialek J, et al. (1997) Coronary risk factors and plaque morphology in men with coronary disease who died suddenly. *N Engl J Med* 336: 1276–1282.
- Davies MJ (1997) The composition of coronary-artery plaques. *N Engl J Med* 336: 1312–1314.
- Virmani R, Burke AP, Farb A, Kolodgie FD (2006) Pathology of the vulnerable plaque. *J Am Coll Cardiol* 47: C13–18.
- Ridker PM, Buring JE, Cook NR, Rifai N (2003) C-reactive protein, the metabolic syndrome, and risk of incident cardiovascular events: an 8-year follow-up of 14,719 initially healthy American women. *Circulation* 107: 391–397.
- Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, et al. (1998) Prediction of coronary heart disease using risk factor categories. *Circulation* 97: 1837–1847.
- Fayad ZA, Nahar T, Fallon JT, Goldman M, Aguinaldo JG, et al. (2000) In vivo magnetic resonance evaluation of atherosclerotic plaques in the human thoracic aorta: a comparison with transesophageal echocardiography. *Circulation* 101: 2503–2509.
- Cappendijk VC, Cleutjens KB, Kessels AG, Heeneman S, Schurink GW, et al. (2005) Assessment of human atherosclerotic carotid plaque components with multisequence MR imaging: initial experience. *Radiology* 234: 487–492.
- Cai JM, Hatsukami TS, Ferguson MS, Small R, Polissar NL, et al. (2002) Classification of human carotid atherosclerotic lesions with in vivo multicontrast magnetic resonance imaging. *Circulation* 106: 1368–1373.
- Fayad ZA, Fuster V, Fallon JT, Jayasundera T, Worthley SG, et al. (2000) Noninvasive in vivo human coronary artery lumen and wall imaging using black-blood magnetic resonance imaging. *Circulation* 102: 506–510.
- Botnar RM, Stuber M, Kissinger KV, Kim WY, Spuentrup E, et al. (2000) Noninvasive coronary vessel wall and plaque imaging with magnetic resonance imaging. *Circulation* 102: 2582–2587.
- Kim WY, Stuber M, Bornert P, Kissinger KV, Manning WJ, et al. (2002) Three-dimensional black-blood cardiac magnetic resonance coronary vessel wall imaging detects positive arterial remodeling in patients with nonsignificant coronary artery disease. *Circulation* 106: 296–299.
- Assmann G, Cullen P, Schulte H (2002) Simple scoring scheme for calculating the risk of acute coronary events based on the 10-year follow-up of the prospective cardiovascular Munster (PROCAM) study. *Circulation* 105: 310–315.
- Botnar RM, Kim WY, Bornert P, Stuber M, Spuentrup E, et al. (2001) 3D coronary vessel wall imaging utilizing a local inversion technique with spiral image acquisition. *Magn Reson Med* 46: 848–854.
- Desai MY, Lai S, Barnet C, Weiss RG, Stuber M (2005) Reproducibility of 3D free-breathing magnetic resonance coronary vessel wall imaging. *Eur Heart J* 26: 2320–2324.
- Zhang S, Hatsukami TS, Polissar NL, Han C, Yuan C (2001) Comparison of carotid vessel wall area measurements using three different contrast-weighted black blood MR imaging techniques. *Magn Reson Imaging* 19: 795–802.
- Spuentrup E, Katoh M, Buecker A, Manning WJ, Schaeffter T, et al. (2004) Free-breathing 3D Steady-State Free Precession Coronary MR Angiography with Radial k-Space Sampling: Comparison with Cartesian k-Space Sampling and Cartesian Gradient-Echo Coronary MR Angiography—Pilot Study. *Radiology* 231: 581–586.
- Kim WY, Astrup AS, Stuber M, Tarnow L, Falk E, et al. (2007) Subclinical coronary and aortic atherosclerosis detected by magnetic resonance imaging in type 1 diabetes with and without diabetic nephropathy. *Circulation* 115: 228–235.
- Adame IM, van der Geest RJ, Wasserman BA, Mohamed MA, Reiber JH, et al. (2004) Automatic segmentation and plaque characterization in atherosclerotic carotid artery MR images. *Magma* 16: 227–234.
- Austen WG, Edwards JE, Frye RL, Gensini GG, Gott VL, et al. (1975) A reporting system on patients evaluated for coronary artery disease. Report of the Ad Hoc Committee for Grading of Coronary Artery Disease, Council on Cardiovascular Surgery, American Heart Association. *Circulation* 51: 5–40.
- O'Brien PC (1984) Procedures for comparing samples with multiple endpoints. *Biometrics* 40: 1079–1087.
- Läuter J (1996) Exact T and F tests for analyzing studies with multiple endpoints. *Biometrics* 52: 964–970.
- Tuzcu EM, Kapadia SR, Tutar E, Ziada KM, Hobbs RE, et al. (2001) High prevalence of coronary atherosclerosis in asymptomatic teenagers and young adults: evidence from intravascular ultrasound. *Circulation* 103: 2705–2710.
- Miao C, Chen S, Macedo R, Lai S, Liu K, et al. (2009) Positive remodeling of the coronary arteries detected by magnetic resonance imaging in an asymptomatic population: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol* 53: 1708–1715.
- Fuster V (1994) Lewis A. Conner Memorial Lecture. Mechanisms leading to myocardial infarction: insights from studies of vascular biology. *Circulation* 90: 2126–2146.
- Strong JP, Mc GH, Jr. (1962) The natural history of coronary atherosclerosis. *Am J Pathol* 40: 37–49.
- Lee RT, Libby P (1997) The unstable atheroma. *Arterioscler Thromb Vasc Biol* 17: 1859–1867.

29. Grundy SM, Balady GJ, Criqui MH, Fletcher G, Greenland P, et al. (1998) Primary prevention of coronary heart disease: guidance from Framingham: a statement for healthcare professionals from the AHA Task Force on Risk Reduction. American Heart Association. *Circulation* 97: 1876–1887.
30. Wood D, De Backer G, Faergeman O, Graham I, Mancia G, et al. (1998) Prevention of coronary heart disease in clinical practice. Recommendations of the Second Joint Task Force of European and other Societies on coronary prevention. *Eur Heart J* 19: 1434–1503.
31. Clevert DA, Johnson T, Michaely H, Jung EM, Flach PM, et al. (2006) High-grade stenoses of the internal carotid artery: comparison of high-resolution contrast enhanced 3D MRA, duplex sonography and power Doppler imaging. *Eur J Radiol* 60: 379–386.
32. Leiner T (2005) Magnetic resonance angiography of abdominal and lower extremity vasculature. *Top Magn Reson Imaging* 16: 21–66.
33. Patel ST, Mills JL, Sr, Tynan-Cuisinier G, Goshima KR, Westerband A, et al. (2005) The limitations of magnetic resonance angiography in the diagnosis of renal artery stenosis: comparative analysis with conventional arteriography. *J Vasc Surg* 41: 462–468.
34. Rumberger JA, Simons DB, Fitzpatrick LA, Sheedy PF, Schwartz RS (1995) Coronary artery calcium area by electron-beam computed tomography and coronary atherosclerotic plaque area. A histopathologic correlative study. *Circulation* 92: 2157–2162.
35. Greenland P, LaBree L, Azen SP, Doherty TM, Detrano RC (2004) Coronary artery calcium score combined with Framingham score for risk prediction in asymptomatic individuals. *Jama* 291: 210–215.
36. de Roos A, Kroft LJ, Bax JJ, Geleijns J (2007) Applications of multislice computed tomography in coronary artery disease. *J Magn Reson Imaging* 26: 14–22.
37. Leber AW, Becker A, Knez A, von Ziegler F, Sirol M, et al. (2006) Accuracy of 64-slice computed tomography to classify and quantify plaque volumes in the proximal coronary system: a comparative study using intravascular ultrasound. *J Am Coll Cardiol* 47: 672–677.
38. Fujii K, Carlier SG, Mintz GS, Wijns W, Colombo A, et al. (2005) Association of plaque characterization by intravascular ultrasound virtual histology and arterial remodeling. *Am J Cardiol* 96: 1476–1483.
39. Schoenhagen P, Ziada KM, Kapadia SR, Crowe TD, Nissen SE, et al. (2000) Extent and direction of arterial remodeling in stable versus unstable coronary syndromes: an intravascular ultrasound study. *Circulation* 101: 598–603.
40. Brenner DJ, Hall EJ (2007) Computed tomography—an increasing source of radiation exposure. *N Engl J Med* 357: 2277–2284.
41. Akhter N, Milford-Beland S, Roe MT, Piana RN, Kao J, et al. (2009) Gender differences among patients with acute coronary syndromes undergoing percutaneous coronary intervention in the American College of Cardiology–National Cardiovascular Data Registry (ACC-NCDR). *Am Heart J* 157: 141–148.
42. Lokhandwala JO, Skelding KA (2009) Sex differences in percutaneous coronary interventions. *J Cardiovasc Transl Res* 2: 300–305.
43. Maintz D, Ozgun M, Hoffmeier A, Fischbach R, Kim WY, et al. (2006) Selective coronary artery plaque visualization and differentiation by contrast-enhanced inversion prepared MRI. *Eur Heart J* 27: 1732–1736.
44. Yeon SB, Sabir A, Clouse M, Martinezclark PO, Peters DC, et al. (2007) Delayed-enhancement cardiovascular magnetic resonance coronary artery wall imaging: comparison with multislice computed tomography and quantitative coronary angiography. *J Am Coll Cardiol* 50: 441–447.
45. Naghavi M, Libby P, Falk E, Casscells SW, Litovsky S, et al. (2003) From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: Part I. *Circulation* 108: 1664–1672.